

Salivary Enhancement: Current status and future therapies

Jane C. Atkinson, D.D.S.

Department of Oral Medicine

University of Maryland Dental School, Baltimore, MD

Bruce J. Baum, D.M.D., Ph.D.²

Gene Therapy and Therapeutics Branch

National Institute of Dental and Craniofacial Research, Bethesda, MD

Correspondence:

Jane C. Atkinson, DDS
University of Maryland, Baltimore College of Dental Surgery
Department of Clinical Affairs
666 West Baltimore Street, Room 3E-32
Baltimore, Maryland 21201-1586
Phone: 301-706-8125
Fax: 301-706-3028
E-Mail: Jatkinson@Dental.umaryland.edu

Abstract:

Saliva provides the principal protective milieu for teeth by modulating oral microbial ecosystems and reversing the initial phases of caries development. Patients with inadequate salivary function are at increased risk for dental decay. Therefore, it is likely that therapies that increase overall fluid output of these individuals will reverse early carious lesions. The most common causes of salivary dysfunction are medication usage, autoimmune exocrinopathy, Sjögren's syndrome, and damage of salivary parenchyma during therapeutic irradiation. For patients with remaining functional acinar tissue, treatment with the parasympathomimetic secretagogues pilocarpine and Cevimeline may provide relief. However, these medications do not benefit all patients. The possibilities of using gene therapy and tissue engineering to develop treatments for those with severe salivary dysfunction are discussed.

Keywords: salivary dysfunction; saliva; xerostomia; Sjögren's syndrome; gene therapy; dry mouth; dental caries

Saliva provides the principal protective milieu for teeth by modulating oral microbial ecosystems, reversing the initial phases of caries development, aiding in the preparation of the food bolus, lubricating oral tissues, and supporting other critical functions.¹ Patients with inadequate salivary function are at increased risk for dental decay, oral and mucosal infections, gastrointestinal complications, and a decreased quality of life.^{2,3}

Patients with significantly decreased salivary output have an increased prevalence of dental caries.^{4,6} Therefore, therapies that increase overall salivary flow in these individuals are believed to have the potential of reversing early carious lesions. Though this has not been established in humans, studies with rats demonstrated that restoration of salivary function with the cholinergic agonist, pilocarpine, decreased new caries formation in animals whose salivary flow was reduced by partial salivary gland removal.⁷

CAUSES OF SALIVARY DYSFUNCTION

Many systemic diseases are associated with alterations in salivary output. The most pronounced salivary dysfunction occurs in patients taking medications that interfere with salivary secretory processes,^{8,9} those who have received therapeutic irradiation to eradicate head and neck malignancies,⁹⁻¹¹ and patients with Sjögren's syndrome (SS; ref. 10-11). Salivary hypofunction secondary to medication usage is likely the most common cause of these conditions.^{8,9}

Medications often inhibit cholinergic signaling pathways in salivary tissues and thereby decrease the fluid output of the gland. Interference with other peripheral and central signaling pathways can also reduce salivary output and alter salivary composition. While 300 to 400 medications are believed to interfere with salivary secretion,^{8,9} the specific inhibitory

mechanisms are defined only for small subsets of drugs. The permanent impact of prolonged anticholinergic medication usage on salivary tissues still requires definition. Often, the only treatment for these patients is to change the medication to a less-xerogenic type or decrease the medication dose below that leading to oral dryness while maintaining the required therapeutic effect. However, this is not possible in many situations. Further, prescription use in the United States likely will continue to increase.^{12,13} At present, elders (> 65 years of age) use an average of about three prescription and two over-the-counter medications per day,^{14,15} and salivary flow decreases as the numbers of medications a patient takes daily increases.¹⁶⁻¹⁷ Given that the number of elders is increasing, we can expect an increase in patients with compromised salivary function secondary to systemic diseases and their treatments. The best solution to this problem would be the development of more specific pharmaceutical agents that selectively act on target tissues (such as vessels) while sparing others, such as salivary glands. In addition, clinical studies are needed to determine if salivary flow rates influence a dental restoration's length of service.

Salivary hypofunction after gland irradiation is very difficult to treat, as salivary parenchyma within the radiation field is damaged permanently.^{18,19} Head and neck cancer affects 30,000 - 40,000 new patients each year, most of whom are treated with therapeutic irradiation. These patients are typically middle-aged males and often individuals from economically disadvantaged backgrounds.^{20,21} Radiation treatment of oral and pharyngeal malignancies typically includes salivary tissue within the field. At doses above 50 Gy, patients can lose all salivary function if the salivary glands are totally within the radiation field.²² The reasons for the extreme radiosensitivity of human salivary tissues remain undefined.^{18,22}

Sjögren's syndrome affects about one million persons in the U.S., currently estimated to reflect a 9:1, female: male ratio.^{23,24} In most patients, the diagnosis is established when the patient is between 40 and 50 years of age. SS is a systemic autoimmune disorder primarily affecting the salivary and lacrimal glands in which clusters of infiltrating lymphocytes replace the parenchyma of the glands.²⁵ During the last decade, many studies have examined the influences of inflammatory proteins on salivary tissue health and function.²⁶⁻²⁷ Autoantibodies were identified in the circulation and saliva of SS patients over forty years ago.²⁸ Passive transfer of these autoantibodies can reduce salivary output in the nonobese diabetic (NOD) mouse model of SS,^{29,30} possibly by blocking cholinergic M3 receptors. Other studies in the last decade have examined autoantibodies to alpha-fodrin³¹ and programmed cell death (apoptosis) in SS.³²⁻³⁵ Still undefined are the events that trigger initial lymphocytic infiltrates and perpetuate the disease.

TREATMENT OF SALIVARY DYSFUNCTION

Both irradiation and Sjögren's syndrome lead to the loss of salivary acinar cells, the only cell type in the glands capable of fluid movement. These conditions exhibit considerable heterogeneity. Some patients experience minimal parenchymal cell loss, while others may have glands entirely replaced by connective tissue and inflammatory cells. Treatment of patients with remaining functional acinar tissue is possible with a parasympathomimetic secretagogue.

The first such drug approved in the U.S. was pilocarpine. Pilocarpine possesses modest, non-specific muscarinic and weak β -adrenergic agonist activity. Its effectiveness in increasing salivary output was demonstrated in several studies of patients with radiation-induced salivary hypofunction and Sjögren's syndrome (Table 1).³⁶⁻⁴⁰ Recently, the FDA approved a second

secretagogue, cevimeline.^{41,42} Cevimeline is a more specific drug, with a preference for activation of the primary muscarinic receptor sub-type responsible for fluid flow from salivary glands, the so-called M3 receptor. This medication has not been tested to date in post-radiation patients. In general, secretory agents do not address either the underlying inflammatory processes of SS or the lack of functional acinar cells after radiation therapy, and will have very limited success in those with advanced salivary dysfunction. Likewise, the interaction of a cholinergic agonist with other medications precludes the use of pilocarpine in many patients with medication-induced salivary dysfunction.

Prevention of salivary damage during radiation therapy should be a goal in oncology. Radiation damage to salivary glands can be limited by pre-radiation planning (conformal and static, multi-segmental intensity modulated technique, IMRT) that spares as much salivary tissue as possible from the radiation field.⁴³ Use of the oxygen radical scavenger amifostine during radiation treatment may decrease damage to glands.⁴⁴ Surgically repositioning of submandibular salivary glands to the submental space before radiation has been used to maintain gland function.⁴⁵

Several anti-inflammatory medications have been tested for the treatment of Sjögren's syndrome (Table 2),⁴⁶⁻⁵² but only interferon alpha and prednisolone irrigation increased salivary output.^{51,52} One reason for this generalized treatment failure is that the doses of medications used in most trials were anti-inflammatory, and not immunosuppressive. The increased risk of serious side effects have discouraged investigators and patients from testing more potent regimens of drugs such as prednisone, cyclosporine and azathioprine. However, new anti-inflammatory and immune-mediating agents are continually being tested as treatments for Sjögren's syndrome.

FUTURE TREATMENTS

There is currently no conventional therapy to enhance salivary secretion for patients with extensive gland damage. This circumstance provided the impetus approximately 10 years ago for the application of gene transfer technology to repair irradiation- or autoimmune-damaged salivary glands. The initial goal of these studies was to re-engineer the function of the surviving non-fluid secreting ductal cells in damaged glands to a secretory phenotype.

The first peer-reviewed publication on gene transfer to salivary glands was published in 1994.⁵³ Since that report, several laboratories have transferred several different genes successfully to salivary glands.⁵⁴ Most of these studies have utilized viral vectors, particularly adenoviral vectors, to mediate gene transfer. Viral vectors typically are extremely efficient at transferring genes, but can pose a safety risk and stimulate potent immune responses. An alternative means of gene transfer is to use non-viral methods. Perhaps the most successful form of non-viral gene transfer to salivary glands involves the use of cationic liposomes.⁵⁵ While much less efficient than viral vectors, liposomes pose relatively little safety risk. Current research is focusing on ways to optimize this method of gene transfer in salivary glands.

In 1997, a study reported by Delporte *et al.*,⁵⁶ described the “correction” of irradiation-induced salivary hypofunction in rats through transfer of the cDNA encoding aquaporin 1, a mammalian water channel (permeability pathway). Gene transfer was accomplished using a replication-deficient, first generation, recombinant adenovirus. Irradiated rats administered a control adenovirus exhibited salivary flow rates approximately 65 percent lower than sham-irradiated animals. Conversely, when animals were administered the aquaporin 1-encoding adenovirus four months after irradiation, salivary flow rates determined three days post-

administration were statistically the same as control levels. The studies were continued in primates⁵⁷ and were not as effective as the Delporte *et al.* rat study. Because of the small number of primates used (five), the reasons for the minimal success are not entirely clear and may have been technical. A larger study is now being conducted in pigs that will eventually lead to a clinical trial in humans.

Thus, the specific value of aquaporin 1 gene transfer for irradiated salivary glands must be considered as hypothetical, and is not ready for clinical testing in humans. It is not known whether the simple insertion of a water channel into surviving ductal cells will correct glandular hypofunction. However, gene transfer without question can be readily accomplished *in vivo* in salivary glands and is potentially of considerable clinical value to enhance salivary secretions.

Gene transfer can also be utilized to augment salivary secretions by transferring genes that encode secretory proteins into salivary glands. The proteins are subsequently secreted in an exocrine manner. This was successfully accomplished in animal studies,⁵⁸ with the transfer of the human histatin 3 cDNA to rat submandibular glands. Histatin 3, which normally is not secreted in rodent saliva, was secreted at high levels (up to 1 mg/ml) after gene transfer. Another potential clinical use of gene transfer to salivary glands is immunization via DNA vaccination. Kawabata *et al*⁵⁹ showed that delivery of the cDNA for the *P. gingivalis* fimbrial protein into murine salivary glands led to the production of secretory immunoglobulin A specific for this microbial protein. This approach could be used to immunize humans against other oral microbes, such as mutans streptococcus. Gene transfer can also be used to deliver anti-inflammatory proteins, such as Interleukin 10, to salivary glands. This may have potential in the treatment of Sjögren's syndrome.⁶⁰

Gene transfer to repair damaged glands can only be an option if epithelial tissue survives either the irradiation or autoimmune damage. In the absence of any parenchymal cells, when a gland is fully replaced by fibrotic tissue, gene transfer cannot lead to an enhancement of saliva production since no system exists to produce and transport fluid into the mouth. Recently, to address this circumstance, we began initial studies directed at the development of an artificial salivary gland using well-established principles of tissue engineering coupled with genetic engineering.⁶¹

The prototype design includes a biodegradable substratum shaped as a blind end tube (i.e., like a test tube), coated with a layer of purified extracellular matrix proteins involved in cellular organization, followed by a monolayer lining of polarized epithelial cells capable of unidirectional fluid secretion.⁶² Initial feasibility studies have been reported.⁶³ Although considerably more work is required, based on the success of other groups with developing functional, complex fluid secreting bioartificial organs, notably bladder,⁶⁴ it is reasonable to expect that an artificial salivary gland suitable for clinical testing can be achieved in approximately 10 years. The most immediate therapy in this area may be the use of autologous salivary tissues harvested from the individual, expanded *ex vivo*, and then re-implanted in an appropriate matrix to induce regrowth and repair. For example, tissue could be harvested prior to a course of head and neck radiotherapy, and placed back into an individual following radiation and a healing period.

We are presently experiencing a time of unparalleled biomedical scientific progress. The human genome has been sequenced, and functions are being identified for these newly discovered genes. The challenge will be managing the great volume of information to make meaningful interpretations in real time that apply to therapy. For example, current European

studies are examining the genetics of SS by studying the presence of SS autoantibodies and different HLA alleles.^{65,66} Understanding the genetic basis of Sjögren's syndrome should lead to earlier and more accurate diagnosis of this condition. Major progress in cellular, developmental and molecular biology also will result in treatments of oral biological problems. Enhancing saliva will be but one of many such examples.

REFERENCES

1. Schenkels LCPM, Veerman ECI, Nieuw Amerongen AV. Biochemical composition of human saliva in relation to other mucosal fluids. *Crit Rev Oral Biol Med* 1995;6:161-75.
2. Eisbruch A, Dawson LA, Kim HM, et al. Conformal and intensity modulated irradiation of head and neck cancer: the potential for improved target irradiation, salivary gland function, and quality of life. *Acta Otorhinolaryngol Belg* 1999;53:271-5.
3. Abraham CM, al-Hashimi I, Haghghat N. Evaluation of the levels of oral Candida in patients with Sjögren's syndrome. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1998;86:65-8.
4. Spak CJ, Johnson G, Ekstrand J. Caries incidence, salivary flow rate and efficacy of fluoride gel treatment in irradiated patients. *Caries Res* 1994;28:388-93.
5. Papas AS, Joshi A, MacDonald SL, Maravelis-Splagounias L, Pretara-Spanedda P, Curro FA. Caries prevalence in xerostomic individuals. *J Can Dent Assoc* 1993;59:171-4, 177-9.
6. Leone CW, Oppenheim FG. Physical and chemical aspects of saliva as indicators of risk for dental caries in humans. *J Dent Educ* 2001;65:xx.
7. O'Connell AC, Pearson SK, Bowen WH. Pilocarpine alters caries development in partially-desalivated rats. *J Dent Res* 1994;73:637-43.
8. Nederfors T. Xerostomia: prevalence and pharmacotherapy. With special reference to beta-adrenoceptor antagonists. *Swed Dent J Suppl.* 1996;116:1-70.
9. Fox PC. Acquired salivary dysfunction. *Drugs and radiation.* *Ann N Y Acad Sci* 1998;842:132-7.
10. Atkinson JC, Fox PC. Salivary gland dysfunction. *Clin Geriatr Med* 1992;8:499-511.
11. Atkinson JC, Wu AJ. Salivary gland dysfunction. *J Am Dent Assoc* 1994;125:409-16.

12. Copeland C. Prescription drugs: issues of cost, coverage, and quality. EBRI Issue Brief 1999; 208:1-21.
13. Briesacher BA, Stuart B, Peluso R. Drug use and prescribing problems in the community-dwelling elderly: a study of three state Medicaid programs. Clin Ther 1999;21:2156-72.
14. Helling KD, Lemke JH, Semla TP, Wallace RB, Lipson DP, Cornoni-Huntley J. Medication use characteristics in the elderly: the Iowa 65+ rural health study. J Am Geriatric Soc 1987;35:4-12.
15. Espino DV, Lichtenstein MJ, Hazuda HP, et al. Correlates of prescription and over-the-counter medication usage among older Mexican Americans: the Hispanic EPESE study. J Am Geriatr Soc 1998;46:1228-34.
16. Ghezzi EM, Wagner-Lange LA, Schork MA, et al. Longitudinal influence of age, menopause, hormone replacement therapy, and other medications on parotid flow rates in healthy women. J Gerontol A Biol Sci Med Sci 2000;55:M34-42.
17. Navazesh M, Brightman VJ, Pogoda JM. Relationship of medical status, medications, and salivary flow rates in adults of different ages. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 1996;81:172-6.
18. Guchelaar HJ, Vermes A, Meerwaldt JH. Radiation-induced xerostomia: pathophysiology, clinical course and supportive treatment. Support Care Cancer 1997;5:281-8.
19. Valdez IH, Atkinson JC, Ship JA, Fox PC. Parotid and submandibular/sublingual function in patients with radiation induced xerostomia: salivary flow rate and sialochemistry. Int J Radiat Oncol Biol Phys 1993;25:41-7.

20. Merlano M, Vitale V, Rosso R, et al. Treatment of advanced squamous-cell carcinoma of the head and neck with alternating chemotherapy and radiotherapy. *N Engl J Med* 1992;327:1115-21.
21. Adelstein DJ, Lavertu P, Saxton JP, et al. Mature results of a phase III randomized trial comparing concurrent chemoradiotherapy with radiation therapy alone in patients with stage III and IV squamous cell carcinoma of the head and neck. *Cancer* 2000;88:876-83.
22. O'Connell AC, Redman RS, Evans RL, Ambudkar IS. Radiation-induced progressive decrease in fluid secretion in rat submandibular glands is related to decreased acinar volume and not impaired calcium signaling. *Radiat Res* 1999;151:150-8.
23. Dafni UG, Tzioufas AG, Staikos P, Skopouli FN, Moutsopoulos HM. Prevalence of Sjögren's syndrome in a closed rural community. *Ann Rheum Dis* 1997;56:521-5.
24. Thomas E, Hay EM, Hajeer A, Silman AJ. Sjögren's syndrome: a community-based study of prevalence and impact. *Br J Rheumatol* 1998;37:1069-76.
25. Fox PC, Speight PM. Current concepts of autoimmune exocrinopathy: immunologic mechanisms in the salivary pathology of Sjögren's syndrome. *Crit Rev Oral Biol Med* 1996;7:144-58.
26. Wu AJ, Kurrasch RH, Katz J, Fox PC, Baum BJ, Atkinson JC. Effect of tumor necrosis factor-alpha and interferon-gamma on the growth of a human salivary gland cell line. *J Cell Physiol* 1994;161:217-26.
27. Atkinson JC, Wu JA. Cytokines in the pathogenesis of Sjögren's syndrome. In: Baum BJ, Cohen MM, eds. *Studies in stomatology and craniofacial biology*. Amsterdam: IOS Press; 1996:409-27.

28. Harley JB, Scofield RH, Reichlin M. Anti-Ro in Sjögren's syndrome and systemic lupus erythematosus. *Rheum Dis Clin North Am* 1992;18:337-58.
29. Esch TR, Taubman MA. Autoantibodies in salivary hypofunction in the NOD mouse. *Ann N Y Acad Sci* 1998;842:221-8.
30. Robinson CP, Brayer J, Yamachika S, et al. Transfer of human serum IgG to nonobese diabetic Igmu null mice reveals a role for autoantibodies in the loss of secretory function of exocrine tissues in Sjögren's syndrome. *Proc Natl Acad Sci U S A* 1998;95:7538-43.
31. Haneji N, Nakamura T, Takio K, et al. Identification of alpha-fodrin as a candidate autoantigen in primary Sjögren's syndrome. *Science* 1997;276:604-7.
32. Fujihara T, Fujita H, Tsubota K, et al. Preferential localization of CD8+ alpha E beta 7+ T cells around acinar epithelial cells with apoptosis in patients with Sjögren's syndrome. *J Immunol* 1999;163:2226-35.
33. Nakamura H, Kawakami A, Tominaga M, et al. Expression of CD40/CD40 ligand and Bcl-2 family proteins in labial salivary glands of patients with Sjögren's syndrome. *Lab Invest* 1999;79:261-9.
34. Fox RI. Sjögren's syndrome. Controversies and progress. *Clin Lab Med* 1997;17:431-44.
35. Casciola-Rosen L, Andrade F, Ulanet D, Wong WB, Rosen A. Cleavage by granzyme B is strongly predictive of autoantigen status: implications for initiation of autoimmunity. *J Exp Med* 1999;190: 815-26.
36. Fox PC, Atkinson JC, Macynski AA, et al. Pilocarpine treatment of salivary gland hypofunction and dry mouth (xerostomia). *Arch Intern Med* 1991;151:1149-52.
37. Johnson JT, Ferretti GA, Nethery WJ, et al. Oral pilocarpine for post-irradiation xerostomia in patients with head and neck cancer. *N Engl J Med* 1993;329:390-5.

38. LeVeque FG, Montgomery M, Potter D, et al. A multicenter, randomized, double-blind, placebo-controlled, dose-titration study of oral pilocarpine for treatment of radiation-induced xerostomia in head and neck cancer patients. *J Clin Oncol* 1993;11:1124-31.
39. Rieke JW, Hafermann MD, Johnson JT, et al. Oral pilocarpine for radiation-induced xerostomia: integrated efficacy and safety results from two prospective randomized clinical trials. *Int J Radiat Oncol Biol Phys* 1995;31:661-9.
40. Vivino FB, Al-Hashimi I, Khan Z, et al. Pilocarpine tablets for the treatment of dry mouth and dry eye symptoms in patients with Sjogren syndrome: a randomized, placebo-controlled, fixed-dose, multicenter trial. P92-01 Study Group. *Arch Intern Med* 1999;159:174-81.
41. Iga Y, Arisawa H, Ogane N, et al. (+/-)-cis-2-methylspiro[1,3-oxathiolane-5,3'-quinuclidine] hydrochloride, hemihydrate (SNI-2011, cevimeline hydrochloride) induces saliva and tear secretions in rats and mice: the role of muscarinic acetylcholine receptors. *Jpn J Pharmacol* 1998;78:373-80.
42. Cevimeline (Evoxac) for dry mouth. *Med Lett Drugs Ther* 2000;42(1084)70.
43. Henson BS, Eisbruch A, D'Hondt E, Ship JA. Two-year longitudinal study of parotid salivary flow rates in head and neck cancer patients receiving unilateral neck parotid-sparing radiotherapy treatment. *Oral Oncol* 1999;35:234-41.
44. Jha N, Seikaly H, McGaw T, Coulter L. Submandibular salivary gland transfer prevents radiation-induced xerostomia. *Int J Radiat Oncol Biol Phys* 2000;46:7-11.
45. Bohuslavizki KH, Klutmann S, Brenner W, et al. Radioprotection of salivary glands by amifostine in high-dose radioiodine treatment. Results of a double-blinded, placebo-

controlled study in patients with differentiated thyroid cancer. *Strahlenther Onkol* 1999;175 (Suppl 4):6-12.

46. Drosos AA, Skopouli FN, Costopoulos JS, Papadimitriou CS, Moutsopoulos HM. Cyclosporin A (CyA) in primary Sjögren's syndrome: a double blind study. *Ann Rheum Dis* 1986;45:732-5.
47. Drosos AA, van Vliet-Dascalopoulou E, Andonopoulos AP, Galanopoulou V, Skopouli FN, Moutsopoulos HM. Nandrolone decanoate (deca-durabolin) in primary Sjögren's syndrome: a double blind pilot study. *Clin Exp Rheumatol* 1988;6:53-7.
48. Kruize AA, Hene RJ, Kallenberg CG, et al. Hydroxychloroquine treatment for primary Sjögren's syndrome: a two year double blind crossover trial. *Ann Rheum Dis* 1993;52:360-4.
49. Fox PC, Datiles M, Atkinson JC, et al. Prednisone and piroxicam for treatment of primary Sjögren's syndrome. *Clin Exp Rheumatol* 1993;11:149-56.
50. Price EJ, Rigby SP, Clancy U, Venables PJ. A double blind placebo controlled trial of azathioprine in the treatment of primary Sjögren's syndrome. *J Rheumatol* 1998;25:896-9.
51. Izumi M, Eguchi K, Nakamura H, Takagi Y, Kawabe Y, Nakamura T. Corticosteroid irrigation of parotid gland for treatment of xerostomia in patients with Sjögren's syndrome. *Ann Rheum Dis* 1998;57:464-9.
52. Ship JA, Fox PC, Michalek JE, Cummins MJ, Richards AB. Treatment of primary Sjögren's syndrome with low-dose natural human interferon-alpha administered by the oral mucosal route: a phase II clinical trial. IFN Protocol Study Group. *J Interferon Cytokine Res* 1999;19:943-51.

53. Mastrangeli A, O'Connell B, Aladib W, Fox PC, Baum BJ, Crystal RG. Direct in vivo adenovirus-mediated gene transfer to salivary glands. *Am J Physiol* 1994;266:G1146-55.
54. Baum BJ, O'Connell BC. In vivo gene transfer to salivary glands. *Crit Rev Oral Biol Med* 1999;10:276-283.
55. Baccaglini L, Hoque ATMS, Wellner RB, et al. Cationic liposome-mediated gene transfer to rat salivary gland epithelial cells in vitro and in vivo. *J Gene Med* (in press).
56. Delporte C, O'Connell BC, He X, et al. Increased fluid secretion after adenoviral-mediated transfer of the aquaporin-1 cDNA to irradiated rat salivary glands. *Proc Natl Acad Sci U S A* 1997;94:3268-73.
57. O'Connell AC, Baccaglini L, Fox PC, et al. Safety and efficacy of adenovirus-mediated transfer of the human aquaporin-1 cDNA to irradiated parotid glands of non-human primates. *Cancer Gene Ther* 1999;6:505-13.
58. O'Connell BC, Xu T, Walsh TJ, et al. Transfer of a gene encoding the anticandidal protein histatin 3 to salivary glands. *Hum Gene Ther* 1996;7:2255-61.
59. Kawabata S, Terao Y, Fujiwara T, Nakagawa I, Hamada S. Targeted salivary gland immunization with plasmid DNA elicits specific salivary immunoglobulin A and G antibodies and serum immunoglobulin G antibodies in mice. *Infect Immun* 1999;67:5863-8.
60. Wang S, Baum BJ, Yamano S, et al. Adenoviral-mediated gene transfer to mouse salivary glands. *J Dent Res* 2000;79:701-8.
61. Baum BJ, Mooney DJ. The impact of tissue engineering on dentistry. *J Am Dent Assoc* 2000;131:309-18.

62. Baum BJ, Wang S, Cukierman E, et al. Re-engineering the functions of a terminally differentiated epithelial cell in vivo. *Ann N Y Acad Sci* 1999;875:294-300.
63. Aframian DJ, Cukierman I, Nikolovski J, Mooney DJ, Yamada KM, Baum BJ. The growth and morphological behavior of salivary epithelial cells on matrix protein-coated biodegradable substrata. *Tissue Engineering* 2000;6:209-16.
64. Oberpenning F, Meng J, Yoo JJ, Atala A. De novo reconstitution of a functional mammalian urinary bladder by tissue engineering. *Nat Biotechnol* 1999;17:149-55.
65. Rischmueller M, Lester S, Chen Z, et al. HLA class II phenotype controls diversification of the autoantibody response in primary Sjögren's syndrome (pSS). *Clin Exp Immunol* 1998;111:365-71.
66. Jonsson R, Nakken B, Halse A-K, Skarstein K, Brokstad K, Haga H-J. Inheritance and immunology in Sjögrens syndrome (in Norwegian). *Tidsskr Nor Lægeforen* 2000;120:811-4.

Table 1. Randomized, placebo-controlled trials evaluating pilocarpine treatment

| Author | Design | Patient diagnosis | Number (n) | Dose | Effect on salivary Flow (statistically) | Comments |
|---------------------|---|---|------------------------------|--|--|--|
| Fox et al.(36) | Placebo 1 month, pilocarpine 5 months | Post-radiation (PR) Sjögren's (SS) Idiopathic xerostomia (IX) | PR = 12 SS = 21 IX = 6 | 5 mg, tid, 5 months | ↑ unstimulated parotid and submandibular flows | |
| Johnson et al. (37) | Placebo vs 5 or 10 mg pilocarpine | PR | 207 | 5 or 10 mg, tid 12 weeks | ↑ unstimulated whole salivary flow rates | Increased incidence of side effects with 10 mg dose |
| Leveque et al. (38) | Placebo vs fixed dose or dose titration | PR | 162 | 2.5 mg, 4 weeks; 5.0 mg, 4 weeks; 10 mg, 4 weeks | ↑ unstimulated whole and parotid salivary flow rates | Best results with continuous treatment > 8 weeks at doses > 2.5 mg tid |
| Rieke et al. (39) | Placebo vs fixed dose or dose titration | PR | 369 | 2.5 mg, 4 weeks; 5.0 mg, 4 weeks; 10 mg, 4 weeks | ↑ unstimulated whole salivary flow rates | Best results with continuous treatment > 8 weeks at doses > 2.5 mg tid |
| Vivino et al. (40) | Placebo vs 2.5 or 5.0 mg pilocarpine | SS | 373 | 2.5 or 5.0 mg qid, 3 months | ↑ unstimulated whole salivary flows | |

Table 2. Salivary flow response of patients with Sjögren's syndrome to treatment with immune modulating agents (controlled trials)

| Treatment | Design | Number (n) | Dose | Effect on salivary Flow (statistically) | Comments |
|-------------------------------|---|---|---|---|--|
| Cyclosporin A (46) | Placebo vs active drug | 10 each group | 5 mg/kg body weight, 6 months | No change | |
| Nandrolone decanoate (47) | Placebo vs active drug | 10 each group | 100 mg biweekly, 6 months | No change | |
| Hydroxy-Chloroquine (48) | 2 yr, cross-over trial (all had drug and placebo) | 19 | 400 mg/day, 12 months | No change | |
| Piroxicam or Prednisone (49) | Placebo vs active drug | 8 each group | 30 mg qod (prednisone) or 20 mg qd (piroxicam) for 6 months | No change | |
| Azathioprine (50) | Placebo vs active drug | 13 azathioprine 12 placebo | 1 mg/kg body weight | No change | 6 patients, all taking azathioprine, withdrew 2° to side effects |
| Prednisolone irrigation (51) | Saline irrigation followed by active treatment | 28 | 2 mg / parotid gland, 1x / week for 3 weeks | ↑ stimulated whole salivary flow | Saline Tx for 8 weeks, then prednisolone Tx |
| Interferon alpha lozenge (52) | Placebo vs four doses of interferon alpha | Placebo = 22; 150 IU* qd = 21 150 IU tid = 22 450 IU qd = 23 450 tid = 21 | 150 IU qd or 150 IU tid or 450 IU qd or 450 tid Tx for 12 weeks | ↑ stimulated whole salivary flow only in the 150 IU tid group | |

* IU = International Units